

**Table 1C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Demographic Indicators** <sup>(2)</sup>

Page: 1  
Run Date: 1 / 10 / 09

Total Records	Number	%	Source of Data	Number	%
Complete Records	26,029	74.5	WIC	34,934	100.0
Prenatal Records	2,824	8.1	Maternal Health	0	0.0
Postpartum Records	6,081	17.4	Other	0	0.0
<b>Total Records Accepted</b>	<b>34,934</b>		<b>Total</b>	<b>34,934</b>	<b>100.0</b>
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%	Age Distribution	Number	%	Education Distribution	Number	%
White, Not Hispanic	18,709	53.6	< 15 Years	115	0.3	< High School	10,190	31.9
Black, Not Hispanic	6,739	19.3	15 - 17 Years	2,290	6.6	High School	14,493	45.4
Hispanic	6,655	19.1	18 - 19 Years	4,838	13.9	> High School	7,243	22.7
American Indian/Alaskan Native	685	2.0	20 - 29 Years	21,598	61.9	<b>Total</b>	<b>31,926</b>	<b>100.0</b>
Asian/Pacific Islander	1,560	4.5	30 - 39 Years	5,671	16.2			
Multiple Races	584	1.7	>= 40 Years	404	1.2			
All Other	0	0.0	<b>Total</b>	<b>34,916</b>	<b>100.0</b>			
<b>Total</b>	<b>34,932</b>	<b>100.0</b>	<i>Excluded Unknown: 18</i>	<i>0.1</i>		<i>Excluded Unknown: 3,008</i>	<i>8.6</i>	
<i>Excluded Unknown: 2</i>	<i>0.0</i>							

% Poverty Level	Number	%	Program Participation <sup>(3)</sup>	Number	%	Migrant Status	Number	%
0 - 50	7,042	24.4	WIC	28,853	100.0	Yes	29	0.1
51 - 100	4,579	15.9	Food Stamps	12,861	29.7	No	34,905	99.9
101 - 130	2,507	8.7	Medicaid	28,559	49.5	<b>Total</b>	<b>34,934</b>	<b>100.0</b>
131 - 150	1,393	4.8	TANF	28,814	1.2			
151 - 185	1,516	5.3	<i>Excluded Unknown:</i>					
186 - 200	2	0.0	WIC:	0	0.0			
> 200	3	0.0	Food Stamps:	15,992	55.4			
Adjunctive Eligibility	11,780	40.9	Medicaid:	294	1.0			
<b>Total</b>	<b>28,822</b>	<b>100.0</b>	TANF:	39	0.1	<i>Excluded Unknown: 0</i>	<i>0.0</i>	
<i>Excluded Unknown: 6,112</i>	<i>17.5</i>							

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	34,504	9.7	1,081,550	11.2
Normal Weight	34,504	43.9	1,081,550	45.1
Overweight	34,504	15.2	1,081,550	15.0
Obese	34,504	31.2	1,081,550	28.6
Maternal Weight Gain (4)				
Less Than Ideal	24,164	24.7	872,976	24.8
Ideal	24,164	30.2	872,976	32.0
Greater Than Ideal	24,164	45.1	872,976	43.1
Anemia (Low Hb/Hct) (5)				
1st Trimester	11,296	7.4	349,143	6.7
2nd Trimester	10,364	11.9	314,271	10.8
3rd Trimester	5,810	33.1	161,665	32.2
Postpartum	29,138	25.8	359,221	29.2
Parity (6)				
0	33,415	40.1	879,923	46.1
1	33,415	28.2	879,923	25.5
2	33,415	17.0	879,923	15.1
≥ 3	33,415	14.6	879,923	13.3
Interpregnancy Interval				
< 6 Months	17,268	12.1	527,136	13.1
6 - <18 Months	17,268	28.8	527,136	26.7
≥18 Months	17,268	59.1	527,136	60.3
Gestational Diabetes	31,161	4.1	150,176	3.6
Hypertension During Pregnancy	31,161	5.6	164,043	3.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI < 19.8), normal weight (BMI 19.9 to 26.0), overweight (BMI >26.0 to 29.0), obese (BMI >29.0).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	31,285	83.3	579,816	78.5
No Care	31,285	2.3	579,816	5.6
WIC Enrollment				
1st Trimester	33,775	33.7	944,971	31.2
2nd Trimester	33,775	31.2	944,971	34.9
3rd Trimester	33,775	17.4	944,971	18.0
Postpartum (3)	33,775	17.7	944,971	15.9
Prepregnancy Multivitamin Consumption (4)	29,810	17.6	38,592	15.7
Multivitamin Consumption During Pregnancy (5)	26,148	83.6	206,967	74.1
Smoking				
3 Months Prior to Pregnancy	33,094	35.6	847,411	27.2
During Pregnancy	27,732	25.7	761,344	16.2
Last 3 Months of Pregnancy	31,575	21.6	692,984	16.8
Postpartum	31,570	24.5	686,323	17.6
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	10,032	33.0	203,249	42.2
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	8,882	28.0	156,923	35.0
Smoking in Household				
During Pregnancy	27,956	22.7	514,582	22.5
Postpartum	31,534	8.8	443,951	24.1
Drinking				
3 Months Prior to Pregnancy	32,154	10.5	785,765	7.4
Last 3 Months of Pregnancy	31,467	0.4	753,112	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Summary of Health Indicators**

Page: 4  
Run Date: 1 / 10 / 09

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	30,255	1.1	972,241	1.1
Low	30,255	6.7	972,241	7.2
Normal	30,255	84.4	972,241	85.0
High	30,255	7.8	972,241	6.8
Preterm (4)	30,697	10.4	873,261	12.1
Full term Low Birth Weight (5)	26,286	3.2	802,123	3.7
Multiple Births	31,941	1.2	935,557	1.5
Breastfeeding				
Ever Breastfed	30,747	65.8	897,379	64.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.

**Table 3C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 5  
Run Date: 1 / 10 / 09

**Comparison of Racial and Ethnic Distribution by Contributor to Nation**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
<b>Wisconsin</b>	<b>34,934</b>	<b>53.6</b>	<b>19.3</b>	<b>19.1</b>	<b>2.0</b>	<b>4.5</b>	<b>1.7</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,153,475</b>	<b>41.5</b>	<b>22.7</b>	<b>28.7</b>	<b>1.1</b>	<b>3.1</b>	<b>0.9</b>	<b>2.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Contributor to Nation**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
<b>Wisconsin</b>	<b>34,934</b>	<b>0.3</b>	<b>6.6</b>	<b>13.8</b>	<b>61.8</b>	<b>16.2</b>	<b>1.2</b>	<b>0.1</b>
<b>Nation (Prior Year)</b>	<b>1,153,475</b>	<b>0.4</b>	<b>6.5</b>	<b>12.6</b>	<b>60.0</b>	<b>19.0</b>	<b>1.4</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Contributor to Nation**

Comparison of Maternal Health Indicators <sup>(2)</sup>														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Wisconsin	34,504	9.7	46.4	24,164	24.7	45.1	5,810	33.1	29,138	25.8	31,161	4.1	31,161	5.6
Nation (Prior Year)	1,081,550	11.2	43.7	872,976	24.8	43.1	161,665	32.2	359,221	29.2	150,176	3.6	164,043	3.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 8  
Run Date: 1 / 10 / 09

**Comparison of Maternal Behavioral Indicators by Contributor to Nation**

Comparison of Maternal Behavioral Indicators <sup>(2)</sup>									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy <sup>(3)</sup>		During Pregnancy <sup>(4)</sup>	
	Number	%	%	Number	%	Number	%	Number	%
Wisconsin	31,285	83.3	2.3	33,775	33.7	29,810	17.6	26,148	83.6
Nation (Prior Year)	579,816	78.5	5.6	944,971	31.2	38,592	15.7	206,967	74.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Contributor to Nation**

Comparison of Smoking Indicators (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%	Number	%
Wisconsin	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
Nation (Prior Year)	847,411	27.2	692,984	16.8	156,923	35.0	514,582	22.5	443,951	24.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Contributor to Nation**

Page: 10  
Run Date: 1 / 10 / 09

<b>Comparison of Infant Health Indicators</b> <sup>(2)</sup>									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
<b>Wisconsin</b>	<b>30,255</b>	<b>7.8</b>	<b>7.8</b>	<b>30,697</b>	<b>10.4</b>	<b>26,286</b>	<b>3.2</b>	<b>30,747</b>	<b>65.8</b>
<b>Nation (Prior Year)</b>	<b>972,241</b>	<b>8.3</b>	<b>6.8</b>	<b>873,261</b>	<b>12.1</b>	<b>802,123</b>	<b>3.7</b>	<b>897,379</b>	<b>64.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 9C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Maternal Health Indicators by Race/Ethnicity, Age, or Education**

Maternal Health Indicators by Race/Ethnicity, Age, or Education														(2)
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity														
White, Not Hispanic	18,522	10.9	44.8	13,343	21.3	48.8	3,130	25.4	15,672	18.7	16,716	3.8	16,716	5.7
Black, Not Hispanic	6,649	8.4	52.7	3,994	27.9	45.3	1,252	54.8	5,530	45.1	5,945	3.3	5,945	7.8
Hispanic	6,528	7.2	48.0	4,965	28.1	38.9	901	29.7	5,616	24.6	5,949	5.8	5,949	4.4
American Indian/Alaskan Native	683	5.7	53.6	473	23.5	47.4	106	37.7	526	32.5	606	5.1	606	3.5
Asian/Pacific Islander	1,543	14.1	27.5	1,012	40.8	23.9	324	31.2	1,329	31.2	1,430	3.2	1,430	1.3
Multiple Races	577	9.9	48.4	377	26.8	49.3	97	*	465	25.4	515	4.5	515	7.0
All Other/Unknown	2	*	*	0	*	*	0	*	0	*	0	*	0	*
Total	34,504	9.7	46.4	24,164	24.7	45.1	5,810	33.1	29,138	25.8	31,161	4.1	31,161	5.6
Age														
< 15 Years	115	17.4	27.0	82	*	*	23	*	106	38.7	111	0.0	111	2.7
15 - 17 Years	2,262	16.3	27.7	1,583	22.2	47.5	370	44.9	1,941	35.4	2,062	1.2	2,062	5.3
18 - 19 Years	4,773	14.6	32.6	3,290	20.9	50.1	733	34.4	4,028	30.8	4,275	1.7	4,275	5.6
20 - 29 Years	21,332	9.4	47.6	15,033	24.6	45.3	3,751	32.8	18,026	25.0	19,327	3.4	19,327	4.9
30 - 39 Years	5,604	4.5	60.6	3,902	28.5	40.2	870	27.9	4,725	20.7	5,035	9.5	5,035	8.0
>= 40 Years	400	4.8	59.8	262	35.9	34.7	62	*	312	20.8	334	14.4	334	12.3
Unknown	18	*	*	12	*	*	1	*	0	*	17	*	17	*
Total	34,504	9.7	46.4	24,164	24.7	45.1	5,810	33.1	29,138	25.8	31,161	4.1	31,161	5.6
Education														
< High School	10,086	11.0	43.6	6,977	27.0	42.5	1,649	39.8	8,526	30.6	9,056	4.0	9,056	4.9
High School	14,357	9.6	47.1	10,199	24.2	46.3	2,477	32.7	12,240	25.3	13,029	3.8	13,029	5.7
> High School	7,169	8.0	49.4	5,163	21.6	48.2	1,255	24.7	6,064	19.5	6,499	4.9	6,499	7.0
Unknown	2,892	10.5	45.3	1,825	28.0	39.9	429	35.0	2,308	26.9	2,577	3.5	2,577	3.9
Total	34,504	9.7	46.4	24,164	24.7	45.1	5,810	33.1	29,138	25.8	31,161	4.1	31,161	5.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 10C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**

**Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education**

<b>Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education</b> (2)									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester	No Care		1st Trimester		Prepregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
<b>Race/Ethnicity</b>									
White, Not Hispanic	16,803	88.0	1.9	18,195	33.6	15,924	21.7	13,671	88.3
Black, Not Hispanic	6,015	78.7	2.5	6,409	30.2	5,843	10.7	5,107	81.3
Hispanic	5,934	79.0	3.0	6,444	39.3	5,662	15.6	5,144	76.5
American Indian/Alaskan Native	598	75.8	2.5	656	39.2	597	11.2	553	85.2
Asian/Pacific Islander	1,411	68.4	3.2	1,502	22.4	1,278	11.0	1,206	68.5
Multiple Races	524	81.5	3.4	569	38.0	506	14.2	466	88.2
All Other/Unknown	0	*	*	0	*	0	*	1	*
<b>Total</b>	<b>31,285</b>	<b>83.3</b>	<b>2.3</b>	<b>33,775</b>	<b>33.7</b>	<b>29,810</b>	<b>17.6</b>	<b>26,148</b>	<b>83.6</b>
<b>Age</b>									
< 15 Years	110	54.5	5.5	105	24.8	105	10.5	84	*
15 - 17 Years	2,065	71.4	3.3	2,203	34.1	1,989	7.1	1,833	81.8
18 - 19 Years	4,312	81.0	2.4	4,619	37.4	4,101	9.7	3,861	81.5
20 - 29 Years	19,420	84.9	2.1	20,905	32.9	18,426	17.7	16,048	84.3
30 - 39 Years	5,031	85.2	2.4	5,529	34.1	4,855	27.3	4,041	84.1
>= 40 Years	330	75.2	2.7	397	29.7	317	32.8	270	87.0
Unknown	17	*	*	17	*	17	*	11	*
<b>Total</b>	<b>31,285</b>	<b>83.3</b>	<b>2.3</b>	<b>33,775</b>	<b>33.7</b>	<b>29,810</b>	<b>17.6</b>	<b>26,148</b>	<b>83.6</b>
<b>Education</b>									
< High School	9,050	76.8	2.8	9,816	36.3	8,713	11.3	8,014	78.5
High School	13,086	84.9	2.2	14,046	34.8	12,488	16.8	11,073	85.2
> High School	6,550	89.7	1.5	7,065	31.0	6,218	27.7	5,130	90.1
Unknown	2,599	81.7	2.9	2,848	26.2	2,391	18.2	1,931	79.0
<b>Total</b>	<b>31,285</b>	<b>83.3</b>	<b>2.3</b>	<b>33,775</b>	<b>33.7</b>	<b>29,810</b>	<b>17.6</b>	<b>26,148</b>	<b>83.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 11C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Smoking Indicators by Race/Ethnicity, Age, or Education**

Smoking Indicators by Race/Ethnicity, Age, or Education <sup>(2)</sup>										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	17,872	48.3	16,939	30.7	6,495	25.7	14,581	26.5	16,933	9.2
Black, Not Hispanic	6,318	24.2	6,023	13.9	1,116	30.4	5,408	27.1	6,015	13.3
Hispanic	6,227	13.4	6,026	6.1	653	45.3	5,576	10.2	6,011	3.8
American Indian/Alaskan Native	648	55.1	612	32.5	284	26.1	580	32.9	605	11.7
Asian/Pacific Islander	1,467	9.9	1,451	5.2	115	47.8	1,312	7.5	1,448	3.9
Multiple Races	561	49.0	524	30.3	219	25.6	498	32.1	522	9.4
All Other/Unknown	1	*	0	*	0	*	1	*	0	*
<b>Total</b>	<b>33,094</b>	<b>35.6</b>	<b>31,575</b>	<b>21.6</b>	<b>8,882</b>	<b>28.0</b>	<b>27,956</b>	<b>22.7</b>	<b>31,534</b>	<b>8.8</b>
Age										
< 15 Years	104	17.3	110	6.4	16	*	93	*	112	24.1
15 - 17 Years	2,151	29.2	2,098	14.7	519	34.3	1,947	36.8	2,094	15.9
18 - 19 Years	4,588	38.9	4,329	20.9	1,441	32.6	4,135	32.6	4,325	12.4
20 - 29 Years	20,416	37.9	19,586	23.3	5,808	27.6	17,140	20.4	19,571	7.8
30 - 39 Years	5,436	27.6	5,096	19.1	1,035	21.2	4,333	16.1	5,081	6.5
>= 40 Years	384	27.6	339	20.4	61	*	297	18.9	334	6.6
Unknown	15	*	17	*	2	*	11	*	17	*
<b>Total</b>	<b>33,094</b>	<b>35.6</b>	<b>31,575</b>	<b>21.6</b>	<b>8,882</b>	<b>28.0</b>	<b>27,956</b>	<b>22.7</b>	<b>31,534</b>	<b>8.8</b>
Education										
< High School	9,621	35.5	9,191	23.6	2,676	22.7	8,606	26.9	9,169	11.7
High School	13,841	40.1	13,206	24.1	4,244	29.5	11,815	24.1	13,192	8.9
> High School	6,926	29.0	6,574	14.9	1,435	35.2	5,439	14.8	6,574	4.8
Unknown	2,706	29.9	2,604	19.2	527	23.1	2,096	18.5	2,599	7.9
<b>Total</b>	<b>33,094</b>	<b>35.6</b>	<b>31,575</b>	<b>21.6</b>	<b>8,882</b>	<b>28.0</b>	<b>27,956</b>	<b>22.7</b>	<b>31,534</b>	<b>8.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 12C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Infant Health Indicators by Race/Ethnicity, Age, or Education**

<b>Infant Health Indicators by Race/Ethnicity, Age, or Education</b> <sup>(2)</sup>									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
<b>Race/Ethnicity</b>									
White, Not Hispanic	16,266	7.1	8.9	16,549	10.0	14,278	2.8	16,505	65.9
Black, Not Hispanic	5,650	12.2	4.2	5,752	13.0	4,674	5.5	5,740	52.3
Hispanic	5,871	5.8	8.8	5,902	9.4	5,170	2.2	6,000	83.0
American Indian/Alaskan Native	572	5.8	14.3	580	11.6	486	3.1	582	59.1
Asian/Pacific Islander	1,398	6.8	3.5	1,406	8.3	1,237	3.5	1,416	49.6
Multiple Races	498	7.8	7.0	508	9.4	441	3.4	504	65.3
All Other/Unknown	0	*	*	0	*	0	*	0	*
<b>Total</b>	<b>30,255</b>	<b>7.8</b>	<b>7.8</b>	<b>30,697</b>	<b>10.4</b>	<b>26,286</b>	<b>3.2</b>	<b>30,747</b>	<b>65.8</b>
<b>Age</b>									
< 15 Years	102	10.8	6.9	102	9.8	85	*	104	51.0
15 - 17 Years	1,999	7.7	5.7	2,031	12.0	1,692	3.5	2,029	57.9
18 - 19 Years	4,161	7.6	6.0	4,185	10.5	3,569	3.8	4,211	64.4
20 - 29 Years	18,809	6.3	7.6	19,061	9.8	16,465	3.0	19,117	65.9
30 - 39 Years	4,844	7.0	10.9	4,977	11.7	4,193	3.1	4,945	70.3
>= 40 Years	323	9.0	10.2	325	14.5	271	4.4	324	66.0
Unknown	17	*	*	16	*	11	*	17	*
<b>Total</b>	<b>30,255</b>	<b>7.8</b>	<b>7.8</b>	<b>30,697</b>	<b>10.4</b>	<b>26,286</b>	<b>3.2</b>	<b>30,747</b>	<b>65.8</b>
<b>Education</b>									
< High School	8,751	7.6	6.4	8,878	11.3	7,501	3.9	8,877	60.9
High School	12,633	6.7	7.6	12,816	10.1	11,051	3.3	12,796	63.8
> High School	6,371	5.4	9.8	6,438	9.8	5,618	2.3	6,490	76.9
Unknown	2,500	6.7	8.7	2,565	10.6	2,116	2.7	2,584	64.9
<b>Total</b>	<b>30,255</b>	<b>7.8</b>	<b>7.8</b>	<b>30,697</b>	<b>10.4</b>	<b>26,286</b>	<b>3.2</b>	<b>30,747</b>	<b>65.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 13C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**

**Maternal Weight Gain and Birth Outcomes by Select Health Indicators**

Maternal Weight Gain and Birth Outcomes by Select Health Indicators <sup>(2)</sup>										
	Weight Gain (3)			Birth Outcomes						
	< Ideal		> Ideal	Birthweight			Preterm (6)		Full Term LBW (7)	
				Low (4)		High (5)				
		Number	%	%	Number	%	%	Number	%	Number
Prepregnancy BMI										
Underweight	2,359	28.6	28.6	2,904	12.3	3.2	2,924	13.9	2,423	5.5
Normal Weight	10,760	25.3	42.2	13,179	7.4	6.2	13,353	10.0	11,539	3.1
Overweight	3,649	14.4	60.0	4,603	7.2	8.3	4,622	9.6	4,027	2.8
Obese	7,366	27.5	47.5	9,343	7.2	11.1	9,432	10.4	8,098	2.8
Weight Gain										
< Ideal	–	–	–	5,584	12.8	4.1	5,675	14.7	4,607	5.1
Ideal	–	–	–	6,986	6.9	5.9	7,012	10.1	6,081	3.1
> Ideal	–	–	–	10,467	5.2	10.9	10,543	7.8	9,403	2.0
Smoking During Pregnancy										
Smokers	4,829	25.0	46.0	6,015	10.3	5.3	6,028	10.8	5,172	5.4
Non-Smokers	14,846	24.6	45.0	18,859	6.3	8.7	18,747	9.6	16,476	2.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 14C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Trends in Racial and Ethnic Distribution**

Page: 16  
Run Date: 1 / 10 / 09

<b>Summary of Trends in Racial and Ethnic Distribution</b> <sup>(2)</sup>								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2007	34,934	53.6	19.3	19.1	2.0	4.5	1.7	0.0
2006	31,077	52.1	18.9	19.5	2.0	4.2	1.6	1.7
2005	29,354	54.7	19.2	17.2	2.2	4.3	0.4	2.0
2004	33,697	57.6	19.4	16.4	2.2	4.2	0.0	0.2
2003	0	*	*	*	*	*	*	*
2002	28,924	58.6	19.7	15.7	2.2	3.8	0.0	0.0
2001	31,080	58.3	20.0	15.5	2.2	4.0	0.0	0.0
2000	38,155	59.3	20.8	13.6	2.1	4.2	0.0	0.0
1999	30,872	61.1	20.9	11.9	2.3	3.9	0.0	0.0
1998	30,319	61.9	21.4	10.5	2.1	4.1	0.0	0.0
1997	30,380	62.3	21.4	9.5	2.4	4.3	0.0	0.0
1996	12,251	84.5	4.2	5.3	1.0	5.1	0.0	0.0
1995	29,036	62.4	21.8	9.0	2.1	4.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 15C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Trends in Age Distribution**

Page: 17  
Run Date: 1 / 10 / 09

<b>Summary of Trends in Age Distribution</b> <sup>(2)</sup>								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2007	34,934	0.3	6.6	13.8	61.8	16.2	1.2	0.1
2006	31,077	0.4	6.7	14.1	62.1	15.5	1.1	0.1
2005	29,354	0.5	6.7	14.0	62.0	15.7	1.2	0.0
2004	33,697	0.5	6.7	14.7	61.0	16.0	1.2	0.0
2003	0	*	*	*	*	*	*	*
2002	28,924	0.5	7.4	15.9	59.6	15.5	1.1	0.0
2001	31,080	0.3	6.6	14.6	60.6	16.8	1.2	0.0
2000	38,155	0.5	8.0	16.4	58.7	15.3	1.1	0.0
1999	30,872	0.6	8.6	16.8	57.7	15.3	1.1	0.0
1998	30,319	0.7	8.8	16.6	57.2	15.6	1.0	0.0
1997	30,380	0.8	9.5	16.0	56.8	15.9	1.0	0.0
1996	12,251	0.4	7.2	15.7	58.7	17.0	1.0	0.0
1995	29,036	1.0	9.7	15.9	56.2	16.1	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 16C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Trends in Maternal Health Indicators**

Summary of Trends in Maternal Health Indicators <sup>(2)</sup>										
Year	Pregpregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2007	34,504	9.7	46.4	24,164	24.7	45.1	5,810	33.1	29,138	25.8
2006	30,755	10.3	46.3	21,643	25.4	44.7	4,588	29.4	24,349	24.9
2005	25,611	10.9	44.0	21,738	27.9	39.9	4,675	30.4	19,010	25.3
2004	29,570	10.9	43.3	26,288	26.9	40.3	5,082	33.2	16,237	26.1
2003	0	*	*	0	*	*	0	*	0	*
2002	25,042	12.2	41.7	21,801	26.0	28.6	4,619	31.0	13,207	25.7
2001	27,250	12.6	40.5	23,624	27.0	39.9	2,855	17.0	14,327	26.4
2000	32,922	13.3	39.3	28,368	27.6	39.2	5,363	26.1	14,148	26.4
1999	27,059	13.8	38.6	23,105	28.3	38.4	4,840	24.5	10,218	25.6
1998	26,850	13.7	38.1	23,658	27.6	38.2	4,515	29.7	9,529	23.2
1997	26,106	13.9	36.8	23,150	28.2	37.9	4,565	27.7	9,599	21.4
1996	10,183	14.9	35.1	9,018	29.2	36.5	1,775	22.3	3,636	18.2
1995	24,837	15.4	34.4	21,746	29.1	36.6	4,377	31.9	8,906	21.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 17C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Trends in Maternal Behavioral Indicators**

Page: 19  
Run Date: 1 / 10 / 09

<b>Summary of Trends in Maternal Behavioral Indicators</b> <sup>(2)</sup>					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2007	31,285	83.3	2.3	33,775	33.7
2006	11,903	83.9	0.1	27,099	29.8
2005	22,622	74.9	12.5	28,229	33.2
2004	30,590	74.3	13.9	32,024	34.7
2003	0	*	*	0	*
2002	25,779	74.1	12.5	27,533	33.2
2001	27,652	74.3	10.7	30,274	0.0
2000	33,694	72.5	12.4	36,735	21.1
1999	26,697	71.1	13.0	29,540	23.2
1998	26,953	71.1	12.4	28,767	36.1
1997	27,168	71.3	12.2	28,118	34.4
1996	10,761	72.9	12.9	11,274	36.0
1995	25,376	70.1	13.3	25,878	32.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 18C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Summary of Trends in Smoking Indicators**

Page: 20  
Run Date: 1 / 10 / 09

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2007	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
2006	20,837	33.1	28,206	21.2	6,146	26.5	14,138	25.1	20,724	11.0
2005	18,399	39.8	22,131	26.4	6,285	30.0	18,584	39.8	26,480	33.6
2004	26,351	40.8	29,526	23.7	9,094	29.3	26,414	41.6	30,700	39.6
2003	0	*	0	*	0	*	0	*	0	*
2002	22,548	42.2	24,910	24.5	6,872	32.9	22,642	43.4	25,745	41.5
2001	23,803	42.5	26,637	25.5	8,350	26.8	23,769	44.9	27,572	42.9
2000	30,624	42.8	31,613	25.6	10,461	27.6	30,764	46.6	33,784	44.9
1999	24,100	44.4	25,437	26.2	8,424	28.5	24,204	47.1	27,080	45.2
1998	29,369	43.2	6,139	25.4	212	26.9	24,115	47.0	6,326	42.3
1997	29,464	42.2	6,059	25.4	0	*	23,678	47.4	6,332	40.4
1996	11,796	45.3	2,505	27.3	0	*	9,450	51.1	2,622	43.1
1995	27,869	41.5	6,453	26.2	0	*	21,792	49.0	6,828	43.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 19C**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Summary of Trends in Infant Health Indicators**

Summary of Trends in Infant Health Indicators <sup>(2)</sup>									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2007	30,255	7.8	7.8	30,697	10.4	26,286	3.2	30,747	65.8
2006	26,732	7.7	7.7	23,522	10.8	20,880	3.0	27,572	64.1
2005	24,726	7.8	8.2	24,324	11.1	21,339	2.9	24,996	62.3
2004	28,209	7.6	8.6	27,611	11.4	24,125	3.0	28,189	62.2
2003	0	*	*	0	*	0	*	0	*
2002	23,758	7.4	9.2	23,341	10.7	20,350	3.1	23,977	59.2
2001	25,501	7.4	9.2	12,367	27.9	8,615	8.5	25,708	57.2
2000	29,964	7.3	9.4	23,960	12.8	20,250	3.9	30,279	55.3
1999	24,383	7.5	9.4	20,706	15.5	17,201	3.6	23,396	51.1
1998	25,133	7.6	9.3	24,536	11.6	21,191	2.8	20,339	42.3
1997	24,451	7.6	9.5	24,093	11.2	20,374	3.1	21,352	43.2
1996	9,702	6.4	11.0	9,469	10.3	8,208	2.5	9,966	56.0
1995	23,260	7.3	9.6	21,362	11.2	18,699	3.1	23,216	50.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 22  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2007	18,522	10.9	44.8	13,343	21.3	48.8	3,130	25.4	15,672	18.7
	2006	16,080	11.8	43.9	11,881	21.8	48.6	2,511	22.4	13,019	17.5
	2005	14,097	12.5	41.8	12,341	24.4	43.3	2,574	22.8	10,446	18.5
	2004	17,145	12.2	41.0	15,366	24.2	43.2	2,882	25.7	8,996	18.9
	2003	0	*	*	0	*	*	0	*	0	*
	2002	14,802	13.9	39.6	13,063	24.0	31.5	2,593	22.6	7,489	18.1
	2001	15,952	14.2	37.9	14,057	24.8	42.1	1,547	10.9	7,780	19.0
	2000	19,541	15.2	37.4	17,112	25.9	41.2	3,060	19.7	7,982	18.8
	1999	16,522	15.4	36.3	14,284	26.1	40.1	2,866	18.1	6,029	17.6
	1998	16,623	15.5	35.8	14,853	25.5	40.1	2,634	22.8	5,922	17.0
Black, Not Hispanic	2007	6,649	8.4	52.7	3,994	27.9	45.3	1,252	54.8	5,530	45.1
	2006	5,812	8.9	54.4	3,875	29.3	44.1	902	52.1	4,483	45.2
	2005	5,109	8.8	52.1	4,449	30.3	40.3	965	52.3	3,563	43.4
	2004	5,962	9.0	50.6	5,359	29.9	39.4	1,092	55.3	3,010	46.4
	2003	0	*	*	0	*	*	0	*	0	*
	2002	5,084	9.9	49.0	4,337	27.4	25.0	949	54.9	2,321	44.0
	2001	5,703	10.5	48.1	4,848	29.2	39.0	750	33.2	3,026	42.7
	2000	7,236	10.4	45.9	6,117	28.2	39.3	1,253	42.5	2,815	45.4
	1999	5,957	11.1	46.3	5,081	30.5	37.6	1,125	41.1	2,023	45.1
	1998	5,992	11.1	44.4	5,274	29.7	36.8	1,091	48.3	1,784	40.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 23  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2007	6,528	7.2	48.0	4,965	28.1	38.9	901	29.7	5,616	24.6
	2006	5,981	6.8	48.4	4,206	28.6	38.6	794	26.4	4,916	24.0
	2005	4,225	6.7	44.8	3,452	34.8	30.9	728	26.0	3,610	24.3
	2004	4,534	7.5	45.3	3,912	30.5	34.9	730	31.1	3,117	25.5
	2003	0	*	*	0	*	*	0	*	0	*
	2002	3,658	7.9	42.0	3,097	28.8	24.3	741	29.8	2,650	28.2
	2001	3,898	8.9	42.0	3,251	29.3	35.5	407	13.0	2,711	26.3
	2000	4,104	9.2	40.3	3,383	29.5	34.6	695	24.6	2,507	26.4
	1999	2,961	10.1	38.7	2,415	29.6	36.6	541	23.5	1,499	28.1
	1998	2,633	9.3	40.8	2,178	27.8	35.8	456	28.9	1,262	24.6
American Indian/Alaskan Native	2007	683	5.7	53.6	473	23.5	47.4	106	37.7	526	32.5
	2006	619	5.8	57.4	420	26.4	47.1	67	*	466	25.8
	2005	581	7.4	58.9	513	24.2	44.4	85	*	431	28.3
	2004	672	6.7	56.0	586	24.4	44.4	95	*	379	24.8
	2003	0	*	*	0	*	*	0	*	0	*
	2002	575	8.0	53.6	503	24.1	27.8	81	*	210	35.2
	2001	634	7.3	51.4	555	24.9	46.7	73	*	220	33.6
	2000	712	11.1	47.3	638	24.5	44.0	88	*	240	33.8
	1999	624	8.8	45.0	527	25.4	43.3	106	25.5	219	23.3
	1998	580	9.8	47.9	504	28.4	44.6	71	*	185	28.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 24  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2007	1,543	14.1	27.5	1,012	40.8	23.9	324	31.2	1,329	31.2
	2006	1,309	14.2	26.9	886	39.7	23.7	243	31.3	1,080	32.9
	2005	1,077	16.4	25.7	908	38.8	22.7	260	33.8	858	35.5
	2004	1,205	16.8	27.1	1,064	39.7	21.2	278	31.3	730	33.2
	2003	0	*	*	0	*	*	0	*	0	*
	2002	923	16.1	26.4	801	42.1	17.9	255	29.8	537	37.2
	2001	1,063	16.2	26.9	913	42.1	20.6	78	*	590	36.8
	2000	1,329	15.9	24.2	1,118	46.3	19.3	267	27.0	604	37.1
	1999	995	16.6	26.2	798	51.9	15.9	202	24.8	448	37.9
	1998	1,022	14.8	24.8	849	49.9	15.4	263	22.8	376	34.0
Multiple Races	2007	577	9.9	48.4	377	26.8	49.3	97	*	465	25.4
	2006	488	9.4	45.7	347	24.2	47.0	67	*	385	26.2
	2005	86	*	*	72	*	*	19	*	76	*
	2004	1	*	*	1	*	*	0	*	1	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 25  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2007	2	*	*	0	*	*	0	*	0	*
	2006	466	12.7	43.3	28	*	*	4	*	0	*
	2005	436	14.7	38.3	3	*	*	44	*	26	*
	2004	51	*	*	0	*	*	5	*	4	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
Total All Race/Ethnic Groups	2007	34,504	9.7	46.4	24,164	24.7	45.1	5,810	33.1	29,138	25.8
	2006	30,755	10.3	46.3	21,643	25.4	44.7	4,588	29.4	24,349	24.9
	2005	25,611	10.9	44.0	21,738	27.9	39.9	4,675	30.4	19,010	25.3
	2004	29,570	10.9	43.3	26,288	26.9	40.3	5,082	33.2	16,237	26.1
	2003	0	*	*	0	*	*	0	*	0	*
	2002	25,042	12.2	41.7	21,801	26.0	28.6	4,619	31.0	13,207	25.7
	2001	27,250	12.6	40.5	23,624	27.0	39.9	2,855	17.0	14,327	26.4
	2000	32,922	13.3	39.3	28,368	27.6	39.2	5,363	26.1	14,148	26.4
	1999	27,059	13.8	38.6	23,105	28.3	38.4	4,840	24.5	10,218	25.6
	1998	26,850	13.7	38.1	23,658	27.6	38.2	4,515	29.7	9,529	23.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 26  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2007	16,803	88.0	1.9	18,195	33.6
	2006	6,548	88.8	0.0	14,333	28.6
	2005	12,797	80.5	10.5	15,420	33.9
	2004	17,664	79.9	11.4	18,387	35.6
	2003	0	*	*	0	*
	2002	15,218	80.2	10.0	16,080	34.7
	2001	16,234	80.0	9.1	17,667	0.0
	2000	20,092	78.2	10.7	21,729	22.4
	1999	16,378	76.0	11.7	17,992	24.5
	1998	16,758	75.9	11.4	17,764	37.8
Black, Not Hispanic	2007	6,015	78.7	2.5	6,409	30.2
	2006	2,401	78.6	0.0	5,031	28.3
	2005	4,516	71.7	10.1	5,478	30.6
	2004	5,927	71.7	11.8	6,346	32.3
	2003	0	*	*	0	*
	2002	4,911	70.2	11.4	5,505	29.4
	2001	5,386	70.9	8.5	6,004	0.0
	2000	6,894	68.9	10.8	7,736	18.5
	1999	5,524	68.7	10.6	6,249	20.6
	1998	5,773	67.3	10.8	6,227	32.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 27  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2007	5,934	79.0	3.0	6,444	39.3
	2006	1,904	81.0	0.2	5,269	33.3
	2005	3,751	63.4	22.7	4,852	35.1
	2004	5,040	61.7	24.8	5,239	37.3
	2003	0	*	*	0	*
	2002	4,096	61.2	22.7	4,303	34.9
	2001	4,291	62.8	18.8	4,704	0.0
	2000	4,547	60.1	21.4	4,970	21.8
	1999	3,129	57.2	23.3	3,490	23.0
	1998	2,771	59.3	21.4	3,012	37.9
American Indian/Alaskan Native	2007	598	75.8	2.5	656	39.2
	2006	294	70.7	0.0	519	36.8
	2005	520	78.7	4.8	616	42.4
	2004	660	72.7	10.8	695	44.0
	2003	0	*	*	0	*
	2002	550	75.3	8.9	593	40.6
	2001	596	73.8	8.2	672	0.0
	2000	700	70.4	12.6	758	26.9
	1999	601	72.4	9.5	649	31.0
	1998	546	69.8	10.1	573	44.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 28  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2007	1,411	68.4	3.2	1,502	22.4
	2006	562	66.5	0.0	1,069	18.1
	2005	996	59.0	13.1	1,203	20.2
	2004	1,298	58.3	17.1	1,338	18.7
	2003	0	*	*	0	*
	2002	1,004	53.1	15.0	1,052	18.3
	2001	1,145	52.8	12.9	1,227	0.0
	2000	1,461	50.0	16.0	1,542	11.8
	1999	1,065	48.8	16.4	1,160	13.8
	1998	1,105	48.5	14.9	1,191	19.6
Multiple Races	2007	524	81.5	3.4	569	38.0
	2006	194	82.0	0.0	432	27.8
	2005	42	*	*	102	24.5
	2004	1	*	*	0	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 29  
Run Date: 1 / 10 / 09

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2007	0	*	*	0	*
	2006	0	*	*	446	65.0
	2005	0	*	*	558	44.6
	2004	0	*	*	19	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
Total All Race/Ethnic Groups	2007	31,285	83.3	2.3	33,775	33.7
	2006	11,903	83.9	0.1	27,099	29.8
	2005	22,622	74.9	12.5	28,229	33.2
	2004	30,590	74.3	13.9	32,024	34.7
	2003	0	*	*	0	*
	2002	25,779	74.1	12.5	27,533	33.2
	2001	27,652	74.3	10.7	30,274	0.0
	2000	33,694	72.5	12.4	36,735	21.1
	1999	26,697	71.1	13.0	29,540	23.2
	1998	26,953	71.1	12.4	28,767	36.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>

Page: 30  
Run Date: 1 / 10 / 09

**Wisconsin**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

Summary of Trends in Smoking Indicators by Race/Ethnicity <sup>(2)</sup>											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2007	17,872	48.3	16,939	30.7	6,495	25.7	14,581	26.5	16,933	9.2
	2006	10,856	46.6	15,000	30.3	4,548	24.4	7,370	29.6	11,021	11.6
	2005	10,309	54.4	12,936	35.2	4,844	28.5	10,372	49.5	14,813	41.5
	2004	14,915	54.7	17,033	31.6	6,932	28.0	14,929	51.1	17,671	48.3
	2003	0	*	0	*	0	*	0	*	0	*
	2002	12,939	56.8	14,651	32.3	5,350	32.2	12,961	52.8	15,175	49.9
	2001	13,605	56.4	15,583	33.5	6,407	26.1	13,591	54.5	16,177	51.3
	2000	17,878	56.0	18,658	33.6	8,039	26.5	17,977	55.6	20,094	53.5
	1999	14,464	56.3	15,529	33.1	6,468	27.8	14,519	55.2	16,552	52.8
1998	18,131	53.5	4,227	28.4	178	28.7	14,544	55.2	4,335	46.3	
Black, Not Hispanic	2007	6,318	24.2	6,023	13.9	1,116	30.4	5,408	27.1	6,015	13.3
	2006	3,921	22.6	5,379	14.2	745	27.2	2,598	29.4	3,935	16.3
	2005	3,604	25.8	4,190	18.7	762	27.4	3,668	34.3	5,199	30.0
	2004	5,228	27.6	5,669	17.8	1,187	27.0	5,266	36.8	5,954	35.1
	2003	0	*	0	*	0	*	0	*	0	*
	2002	4,477	28.7	4,741	19.1	866	27.4	4,511	38.4	4,922	36.6
	2001	4,818	30.2	5,151	19.8	1,126	24.1	4,821	40.0	5,366	38.4
	2000	6,485	29.3	6,483	18.4	1,434	26.5	6,535	40.7	6,909	39.0
	1999	5,135	31.2	5,230	19.6	1,180	25.4	5,166	41.0	5,607	39.2
1998	6,347	31.5	1,097	24.4	21	*	5,374	40.1	1,137	39.4	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>

Page: 31  
Run Date: 1 / 10 / 09

**Wisconsin**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

Summary of Trends in Smoking Indicators by Race/Ethnicity <sup>(2)</sup>											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2007	6,227	13.4	6,026	6.1	653	45.3	5,576	10.2	6,011	3.8
	2006	4,269	12.0	5,552	5.3	455	44.0	2,912	12.7	4,051	5.2
	2005	3,228	11.7	3,526	6.1	319	52.7	3,274	17.2	4,610	14.4
	2004	4,483	13.5	4,896	5.7	516	51.2	4,492	18.1	5,077	17.3
	2003	0	*	0	*	0	*	0	*	0	*
	2002	3,722	12.1	3,986	4.8	343	56.0	3,742	20.1	4,081	19.3
	2001	3,771	13.4	4,174	5.9	410	45.9	3,756	21.5	4,276	20.9
	2000	4,237	13.9	4,356	6.2	479	49.5	4,223	23.7	4,577	22.4
	1999	2,897	17.7	3,028	7.7	404	45.0	2,902	25.2	3,203	23.8
1998	3,052	17.1	548	7.1	9	*	2,604	24.8	585	24.4	
American Indian/Alaskan Native	2007	648	55.1	612	32.5	284	26.1	580	32.9	605	11.7
	2006	437	50.8	562	33.8	196	24.5	302	37.4	425	11.3
	2005	453	68.2	509	43.4	271	32.5	455	58.9	578	50.7
	2004	628	65.1	628	41.6	345	26.1	629	61.7	663	58.4
	2003	0	*	0	*	0	*	0	*	0	*
	2002	532	64.3	530	40.6	247	30.8	546	59.2	553	57.9
	2001	582	70.3	584	45.5	341	22.9	588	63.4	604	62.1
	2000	687	69.0	689	41.7	409	26.2	691	63.4	716	61.6
	1999	597	62.6	587	38.7	314	29.6	605	60.7	614	60.7
1998	631	61.6	101	38.6	3	*	537	62.8	102	55.9	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

Page: 32  
Run Date: 1 / 10 / 09

Summary of Trends in Smoking Indicators by Race/Ethnicity <sup>(2)</sup>											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2007	1,467	9.9	1,451	5.2	115	47.8	1,312	7.5	1,448	3.9
	2006	1,002	8.4	1,251	3.8	75	*	714	8.0	945	5.7
	2005	799	12.6	930	6.5	87	*	803	21.0	1,180	17.5
	2004	1,097	12.0	1,299	5.1	114	45.6	1,098	21.1	1,334	20.0
	2003	0	*	0	*	0	*	0	*	0	*
	2002	878	9.7	1,002	4.9	66	*	882	21.3	1,014	20.4
	2001	1,027	8.1	1,145	3.6	66	*	1,013	16.5	1,149	16.5
	2000	1,337	8.7	1,427	4.6	100	37.0	1,338	18.2	1,488	17.9
	1999	1,007	7.2	1,063	2.9	58	*	1,012	15.4	1,104	14.6
	1998	1,208	5.7	166	6.0	1	*	1,056	14.8	167	12.0
Multiple Races	2007	561	49.0	524	30.3	219	25.6	498	32.1	522	9.4
	2006	346	41.9	458	25.3	127	26.0	236	27.1	343	16.3
	2005	6	*	40	*	2	*	12	*	100	22.0
	2004	0	*	1	*	0	*	0	*	1	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

Page: 33  
Run Date: 1 / 10 / 09

Summary of Trends in Smoking Indicators by Race/Ethnicity <sup>(2)</sup>											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2007	1	*	0	*	0	*	1	*	0	*
	2006	6	*	4	*	0	*	6	*	4	*
	2005	0	*	0	*	0	*	0	*	0	*
	2004	0	*	0	*	0	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
1998	0	*	0	*	0	*	0	*	0	*	
Total All Race/Ethnic Groups	2007	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
	2006	20,837	33.1	28,206	21.2	6,146	26.5	14,138	25.1	20,724	11.0
	2005	18,399	39.8	22,131	26.4	6,285	30.0	18,584	39.8	26,480	33.6
	2004	26,351	40.8	29,526	23.7	9,094	29.3	26,414	41.6	30,700	39.6
	2003	0	*	0	*	0	*	0	*	0	*
	2002	22,548	42.2	24,910	24.5	6,872	32.9	22,642	43.4	25,745	41.5
	2001	23,803	42.5	26,637	25.5	8,350	26.8	23,769	44.9	27,572	42.9
	2000	30,624	42.8	31,613	25.6	10,461	27.6	30,764	46.6	33,784	44.9
	1999	24,100	44.4	25,437	26.2	8,424	28.5	24,204	47.1	27,080	45.2
1998	29,369	43.2	6,139	25.4	212	26.9	24,115	47.0	6,326	42.3	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 34  
Run Date: 1 / 10 / 09

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2007	16,266	7.1	8.9	16,549	10.0	14,278	2.8	16,505	65.9
	2006	14,311	7.3	8.7	12,778	10.0	11,460	3.0	14,678	64.1
	2005	13,898	7.1	9.1	13,584	10.4	12,048	2.5	13,937	62.5
	2004	16,293	7.1	9.5	15,806	10.9	13,884	2.8	16,277	62.9
	2003	0	*	*	0	*	0	*	0	*
	2002	14,008	6.7	10.5	13,660	10.1	12,037	2.8	14,083	60.2
	2001	14,948	6.9	10.4	7,333	32.1	4,824	8.0	15,017	59.3
	2000	17,793	6.5	10.8	14,075	12.6	11,983	3.3	17,813	57.8
	1999	14,943	6.6	10.8	12,633	15.3	10,538	3.1	14,288	54.2
	1998	15,593	6.7	10.9	15,129	10.2	13,282	2.3	12,538	46.6
Black, Not Hispanic	2007	5,650	12.2	4.2	5,752	13.0	4,674	5.5	5,740	52.3
	2006	4,937	11.5	4.2	4,262	14.0	3,638	4.7	5,219	49.7
	2005	4,735	11.9	5.0	4,712	14.4	3,982	4.6	4,839	48.6
	2004	5,441	11.3	5.2	5,469	14.6	4,602	4.9	5,390	46.4
	2003	0	*	*	0	*	0	*	0	*
	2002	4,464	11.4	4.8	4,519	13.3	3,760	4.8	4,579	44.1
	2001	4,878	11.1	4.9	2,408	19.7	1,843	11.6	4,971	39.8
	2000	6,017	11.1	5.0	5,036	13.9	4,110	6.0	6,241	41.2
	1999	4,999	11.6	5.0	4,329	16.7	3,521	6.1	4,833	35.0
	1998	5,291	11.5	5.1	5,277	14.6	4,369	5.1	4,531	27.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 35  
Run Date: 1 / 10 / 09

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2007	5,871	5.8	8.8	5,902	9.4	5,170	2.2	6,000	83.0
	2006	5,322	5.7	8.9	4,661	10.3	4,150	1.8	5,453	81.8
	2005	4,299	5.7	9.0	4,217	9.9	3,762	2.5	4,349	81.4
	2004	4,660	5.1	9.3	4,558	9.1	4,098	2.0	4,688	81.4
	2003	0	*	*	0	*	0	*	0	*
	2002	3,817	5.6	9.5	3,726	9.4	3,312	2.7	3,862	77.6
	2001	4,024	5.3	9.9	1,794	26.9	1,272	6.8	4,068	75.5
	2000	4,136	5.9	10.0	3,268	12.9	2,778	3.4	4,211	72.9
	1999	2,872	5.7	10.4	2,390	14.7	2,007	2.4	2,744	68.5
	1998	2,626	5.4	8.0	2,565	12.1	2,206	1.8	1,913	57.2
American Indian/Alaskan Native	2007	572	5.8	14.3	580	11.6	486	3.1	582	59.1
	2006	534	6.2	12.9	452	9.5	406	1.7	549	65.8
	2005	540	6.3	16.5	531	11.5	464	1.9	532	62.0
	2004	588	5.4	14.3	563	13.5	477	1.5	577	60.8
	2003	0	*	*	0	*	0	*	0	*
	2002	509	4.1	15.5	480	11.7	418	1.2	481	61.5
	2001	539	4.3	16.7	221	19.5	170	7.1	525	59.2
	2000	641	4.2	16.5	465	12.3	397	2.8	624	56.7
	1999	543	4.4	17.1	441	12.5	381	1.6	520	56.5
	1998	515	8.2	17.3	465	12.9	399	2.0	396	44.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

Page: 36  
Run Date: 1 / 10 / 09

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2007	1,398	6.8	3.5	1,406	8.3	1,237	3.5	1,416	49.6
	2006	1,189	5.9	4.7	982	8.1	896	2.7	1,218	44.8
	2005	1,141	7.2	4.3	1,104	9.6	991	3.2	1,148	45.6
	2004	1,226	7.0	5.6	1,215	9.9	1,064	3.0	1,254	49.5
	2003	0	*	*	0	*	0	*	0	*
	2002	960	7.3	5.1	956	11.9	823	3.9	972	42.2
	2001	1,112	5.6	5.2	611	16.4	506	7.3	1,127	38.6
	2000	1,377	6.3	5.2	1,116	10.8	982	3.9	1,390	32.7
	1999	1,026	7.9	4.5	913	16.1	754	4.4	1,011	33.1
	1998	1,108	6.1	4.9	1,100	13.3	935	2.4	961	23.4
Multiple Races	2007	498	7.8	7.0	508	9.4	441	3.4	504	65.3
	2006	439	10.0	5.9	387	14.5	330	4.2	452	64.6
	2005	95	*	*	93	*	81	*	100	68.0
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>

Page: 37  
Run Date: 1 / 10 / 09

Wisconsin

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2007	0	*	*	0	*	0	*	0	*
	2006	0	*	*	0	*	0	*	3	*
	2005	18	*	*	83	*	11	*	91	*
	2004	1	*	*	0	*	0	*	3	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2007	30,255	7.8	7.8	30,697	10.4	26,286	3.2	30,747	65.8
	2006	26,732	7.7	7.7	23,522	10.8	20,880	3.0	27,572	64.1
	2005	24,726	7.8	8.2	24,324	11.1	21,339	2.9	24,996	62.3
	2004	28,209	7.6	8.6	27,611	11.4	24,125	3.0	28,189	62.2
	2003	0	*	*	0	*	0	*	0	*
	2002	23,758	7.4	9.2	23,341	10.7	20,350	3.1	23,977	59.2
	2001	25,501	7.4	9.2	12,367	27.9	8,615	8.5	25,708	57.2
	2000	29,964	7.3	9.4	23,960	12.8	20,250	3.9	30,279	55.3
	1999	24,383	7.5	9.4	20,706	15.5	17,201	3.6	23,396	51.1
	1998	25,133	7.6	9.3	24,536	11.6	21,191	2.8	20,339	42.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.